

Letter to the Editor

Annotations on Indeterminate Cytology of Thyroid Nodules in Thyroidology: Novi Sub Sole?

Check for updates

Keywords: Thyroid gland Guideline Indeterminate cytology TI-RADS Endocrine Surgery Endocrine Pathology

Received: April 14, 2025 Accepted: May 04, 2025 First Published: May 10, 2025

Copyright: © 2025 Sengul. This is an open access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Sengul I, Sengul D. Annotations on indeterminate cytology of thyroid nodules in thyroidology: Novi sub sole?. Barw Medical Journal. 2025;3(3) https://doi.org/10.58742/bmj.v3i3.174

Dear Editor,

Indeterminate cytology (IC) remains the most challenging issue for health professionals working in thyroidology, thyroidologists [1-4]. We read a great deal of the article by Ali *et al* [5]. entitled "Clinicopathological Features of Indeterminate Thyroid Nodules: A Single-center Cross-sectional Study," published in 3rd volume, *Barw Medical Journal*. This study addresses a challenging and crucial issue by examining the characteristics and malignancy rates of thyroid nodules with IC, the most controversial category for The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC). The authors evaluated the clinicopathological features of the thyroid nodules with Category III, TBSRTC, in a single-center cross-sectional study [5].

One of the strengths of the article is its focus on the challenges in managing IC. Ali and colleagues [5] thoroughly examine comprehensive data, including demographic details, medical history, laboratory tests, preoperative imaging, cytologic evaluation, and histopathological diagnosis. The results indicate a notable malignancy rate in Category III, TBSRTC. Furthermore, the study points out that malignancy tended to be younger, while benign nodules were significantly larger than malignant ones. The study also found a significant association between malignant nodules and Thyroid Imaging Reporting and Data System (TI-RADS) categories 4 and 5 and benign with TI-RADS 2 and 3, which findings align with some existing literature, providing valuable insights into the clinical assessment of IC.

However, several limitations of the study warrant consideration. Firstly, its singlecenter and retrospective design may limit the generalizability of the findings to diverse populations and settings. As the authors acknowledge, the retrospective data collection might have resulted in missing crucial information. While TI-RADS scoring was provided, more specific ultrasound features of thyroid nodules could have been beneficial. Of note, does including or excluding noninvasive follicular thyroid neoplasm with papillary-like nuclear features (NIFTP), which has been considered a low-risk entity by the current understanding, affect and/or alter the overall results and the assessment of diagnostic performance and study outcome(s)? [2-4] Furthermore, which caliber of the needle had been utilized throughout the study with or without local and/or topical anesthetic agent(s), and would the utilization of thicker or finer needles in order to obtain cytologic samples with or without any local and/or topical anesthesia alter the outcome(s) of this study? [2] Moreover, which edition of TBSRTC has been used for the work and would stress the up-to-date 3rd edition of TBSRTC [3], considering both the novel and crucial subdivisions of category III might affect the study's relevant outcome(s)? [3,4] Another point of attention is the relatively short data collection period compared to the publication. Finally, while the discussion section compares the findings with various studies in the literature, a more in-depth exploration of the methodological differences and potential discrepancies in results could have been provided. For instance, the conflicting views in the literature regarding the relationship between nodule size and malignancy risk could have been further contrasted with the study's findings. The authors also acknowledge the small sample size as a limitation. For future research, multi-center and prospective studies with detailed imaging, such as elastography and contrast-enhanced sonography, and investigations into the role of molecular markers in thyroid nodules with Category III could improve diagnostic accuracy and potentially reduce unnecessary surgical interventions.

In conclusion, this study significantly contributes to the evaluation of IC in thyroidology despite its limitations. However, considering the noted limitations, further research with more comprehensive and methodologically robust studies in this area is warranted. This issue merits further investigation.

Sincerely.

Ilker Sengul^{*} Division of Endocrine Surgery, Faculty of Medicine, Giresun University, Giresun, Turkey Department of General Surgery, Faculty of Medicine, Giresun University, Giresun, Turkey Demet Sengul Department of Pathology, Faculty of Medicine, Giresun University, Giresun, Turkey * Corresponding author: <u>ilker.sengul.52@gmail.com</u> (I. Sengul). TR28100 Giresun, Turkey

Conflicts of interest: The authors have no conflicts of interest to disclose.

References

- Shishani K, Nawafleh H, Jarrah S, Froelicher ES. Smoking patterns among Jordanian health professionals: a study about the impediments to tobacco control in Jordan. Eur J Cardiovasc Nurs. 2011;10(4):221-7. doi:10.1016/j.ejcnurse.2010.08.001
- Sengul I, Sengul D. Proposal of a novel terminology: minimally invasive FNA and thyroid minimally invasive FNA; MIFNA and thyroid MIFNA. Ann Ital Chir. 2021;92:330-1. <u>doi:10.1590/1806-9282.20210922</u>
- Sengul I, Sengul D. Blurred lines for management of thyroid nodules in the era of AUS/FLUS: Novel subdivisions of Category IIIA and IIIB in a possible forthcoming TBSRTC, 3rd ed.; amending vs. unnecessary? Rev Assoc Med Bras 2021;67(10):1385-6. <u>doi:10.1590/1806-9282.20210763</u>
- Sengul I, Sengul D. Evangely, the subcategorization has been announced in the 2023 Bethesda system for reporting thyroid cytopathology: let bygones be bygones in thyroidology! Rev Assoc Med Bras 2024;70(3):e20231511. doi:10.1590/1806-9282.20231511
- Ali R, Salih AM, Abdullah HO, Abdullah AM, Ali RM, Qaradakhy AJ, et al. Clinicopathological Features of Indeterminate Thyroid Nodules: A Single-center Cross-sectional Study. Barw Med J. 2024;2(4):8-14. <u>doi:10.58742/bmj.v2i4.137</u>